

Aureomycin[®]

100 Granular



**Chlortetracycline
Type A Medicated Article**

See mixing directions, claims, cautions
and warnings on back

Net wt 50 LB (22.68 kg)

 **ALPHARMA.**

Aureomycin[®]
100 Granular

Aureomycin[®]

100 Granular

Chlortetracycline

Type A Medicated Article

Active drug ingredient	Chlortetracycline calcium complex equivalent to 100 g chlortetracycline hydrochloride per lb.													
Ingredients	Dried <i>Streptomyces aureofaciens</i> Fermentation Product and Calcium Sulfate.													
For use in the manufacture of medicated animal feeds. For use in dry feed only. Not for use in liquid medicated feeds.														
Use directions	Mix sufficient Aureomycin 100 Granular Medicated Article to supply desired concentration of chlortetracycline per ton with part of the feed ingredients to make a preblend. Add the remainder of the ingredients and mix thoroughly. For specific use levels, see Indications.													
Mixing directions	<table border="1"> <thead> <tr> <th>Level desired grams per ton</th> <th>Amount of medicated article per ton[†]</th> </tr> </thead> <tbody> <tr> <td>50</td> <td>1/2 lb</td> </tr> <tr> <td>100</td> <td>1 lb</td> </tr> <tr> <td>200</td> <td>2 lb</td> </tr> <tr> <td>400</td> <td>4 lb</td> </tr> <tr> <td>800</td> <td>8 lb</td> </tr> </tbody> </table>	Level desired grams per ton	Amount of medicated article per ton [†]	50	1/2 lb	100	1 lb	200	2 lb	400	4 lb	800	8 lb	<p>† It is recommended that 1 pound of Aureomycin 100 Granular Type A Medicated Article be diluted with 3 pounds of one of the feed ingredients to form a 4 pound working preblend. Use 2 pounds of the working preblend to make a preblend (see Use directions) for a Type C feed containing 50 g Aureomycin chlortetracycline / ton of feed.</p>
Level desired grams per ton	Amount of medicated article per ton [†]													
50	1/2 lb													
100	1 lb													
200	2 lb													
400	4 lb													
800	8 lb													

Indications	Chlortetracycline mg per lb daily	Indications	Chlortetracycline g per ton
Cattle Calves (up to 250 lb): Increased rate of weight gain and improved feed efficiency. Beef Cattle (over 700 lb): Control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline. Beef and Non-Lactating Dairy Cattle: As an aid in control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline when delivered in a free-choice feed. Free-choice feed must be manufactured under a feed mill license utilizing an FDA approved formulation. Calves, Beef and Non-Lactating Dairy Cattle. Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> organisms susceptible to chlortetracycline. Feed for not more than 5 days. The appropriate amount of Aureomycin-containing feed supplement may be mixed in the cattle's daily ration or administered as a top-dress. If the Aureomycin-containing feed supplement is administered as a top-dress, it must be spread uniformly on top of the ration and sufficient space must be provided so that all cattle can eat at the same time.	0.1 0.5 0.5-2.0 10	Swine Increased rate of weight gain and improved feed efficiency. Reduction in the incidence of cervical lymphadenitis (fowl abscesses) caused by Group E Streptococci susceptible to chlortetracycline. Breeding Swine: Control of leptospirosis (reducing the incidence of abortion and shedding of leptospires) caused by <i>Leptospira pomona</i> susceptible to chlortetracycline. Feed continuously for not more than 14 days. Sheep Increased rate of weight gain and improved feed efficiency. Ducks Control and treatment of fowl cholera caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline. Feed in complete ration to provide from 8 to 28 mg per pound of body weight per day depending upon age and severity of disease. Feed for not more than 21 days.	10-50 50-100 400 20-50 200-400
Swine Control of porcine proliferative enteropathies (peltis) caused by <i>Lawsonia intracellularis</i> susceptible to chlortetracycline. Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline. (Note: this drug level is equivalent to approximately 400 grams per ton, depending on feed consumption and body weight.) Feed for not more than 14 days.	10	Chickens Increased rate of weight gain and improved feed efficiency. Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline. Feed continuously for 7 to 14 days. Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Erythrorhynchus coli</i> susceptible to chlortetracycline. Feed continuously for 7 to 14 days. Reduction of mortality due to <i>Escherichia coli</i> infections susceptible to chlortetracycline. Feed for 5 days.	10-50 100-200 200-400 500
Turkeys Control of complicating bacterial organisms associated with hemorrhagic enteritis, enteric enteritis susceptible to chlortetracycline. Feed continuously for 7 to 14 days.	25	Turkeys Increased rate of weight gain and improved feed efficiency. Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline. Feed continuously for 7 to 14 days. Control of hexamitiasis caused by <i>Hexamita meleagridis</i> susceptible to chlortetracycline. Feed continuously for 7 to 14 days. Turkey Poults not over 4 weeks of age: Reduction of mortality due to peratyphoid caused by <i>Salmonella typhimurium</i> susceptible to chlortetracycline.	10-50 200 400 400
Indications	mg per head per day	Indications	mg per g head
Cattle Calves (250 to 400 lb): Increased rate of weight gain and improved feed efficiency. Growing Cattle (over 400 lb): Increased rate of weight gain, improved feed efficiency, and reduction of liver condemnation due to liver abscesses. Beef Cattle: Control of bacterial pneumonia associated with shipping fever complex caused by <i>Pasteurella sp.</i> susceptible to chlortetracycline. Beef Cattle (under 700 lb): Control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline.	25-70 70 350 350	Poultry birds Warning: Peltitosis, or avian chlamydia, or ornithosis is a reportable communicable disease, transmissible between wild and domestic birds, other animals and man. Contact appropriate public health and regulatory officials. Caution: Aspergilliosis may occur following prolonged treatment. Treatment of peltitine birds (parrots, macaws, cockatoos) suspected or known to be infected with peltitosis caused by <i>Chlamydia psittaci</i> sensitive to chlortetracycline. Feed continuously for 45 days. Each bird should consume an amount of medicated feed equal to one-third of its body weight daily. During treatment, parrots, macaws, and cockatoos should be kept individually or in pairs in clean cages.	10
Sheep Breeding Sheep: Reduction in the incidence of (vibriosis) abortion caused by <i>Campylobacter fetus</i> infection susceptible to chlortetracycline.	50		

Warning A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Do not feed to ducks or turkeys producing eggs for human consumption.

NADA 48-751, Approved by FDA

Manufactured by **ALPHARMA**
Alpharma Inc.
Bridgeview, New Jersey 07007

Take Time
Observe Label Directions
*Reference registered by Alpharma Inc.

Net wt 50 LB (22.68 kg)

700385 0902