



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the US product label or package insert.

COCCIVAC* -T (TCV-4M5)

Schering-Plough

Coccidiosis Vaccine

Live Oocysts

DIRECTIONS FOR USE

READ FULL DIRECTIONS BELOW CAREFULLY

Do not vaccinate within 21 days of slaughter.

Store vaccine in refrigerator under 45°F (7°C).

Use entire contents of each vial when first opened.

This vaccine contains gentamicin as a preservative.

CAUTION: Burn containers and all unused contents.

DESCRIPTION

This product contains live oocysts of the following species of coccidia: *E. adenoides*, *E. meleagrimitis*, *E. gallopavonis*, and *E. dispersa*, to aid in the prevention of coccidiosis in turkeys.

INDICATIONS FOR USE

For vaccination of healthy turkeys at one day of age by spray cabinet administration to aid in the prevention of coccidiosis in turkeys.

VACCINATION PROGRAMS

Many factors must be considered in determining the vaccination program for a particular farm or poultry operation. To be fully effective, the vaccine must be administered to healthy receptive birds held in proper environment under good management. In addition, the response may be modified by the age of the birds and their immune status. Seldom does one vaccination under field conditions produce complete protection for all individuals in a given flock. The amount of protection required will vary with the type of operation and the degree of exposure that a flock is likely to encounter. For these reasons a program of periodic revaccination may be required.

PRECAUTIONS

ONLY VACCINATE HEALTHY BIRDS

Consult your poultry pathologist for further recommendations based on conditions existing in your area at any given time.

This product is not ordinarily recommended for use with prestarter or starter feeds containing coccidiostats.

Poults must have access to their droppings as reinfection is required to induce immunity.

SPRAY CABINET ADMINISTRATION

FOR POULTS 1 DAY OF AGE

The vaccine should be diluted with 280 ml of distilled water per 1000 doses of vaccine. Each box of 100 poults should receive 28 ml of vaccine solution. (Dye may be added as a marker.)

Full Directions for use of the Spray Cabinet are available from the Company.

NOTICE

All Schering-Plough Animal Health vaccines released for sale meet the requirements of the licensing authority (U.S. Department of Agriculture) in regard to safety, purity, potency and the capacity to immunize normal susceptible birds.

CAUTION

DO NOT FREEZE

The capacity of this vaccine to produce satisfactory results depends on many factors including, but not limited to conditions of storage and handling by the user, administration of the vaccine, health and responsiveness of individual animals and degree of field exposure. Therefore, directions for use should be followed carefully.

For Veterinary Use Only

U.S. FEDERAL REGULATIONS PROHIBIT THE REPACKAGING OR SALE OF THE CONTENTS OF THIS CARTON IN FRACTIONAL UNITS. DO NOT ACCEPT IF SEAL IS BROKEN.

U.S. Veterinary License No. 226

Schering-Plough Animal Health Corp., Millsboro, DE 19966 U.S.A.

1-800-219-9286 (U.S.A.)

Copyright © 2006