

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

# DANGER

## KEEP OUT OF REACH OF CHILDREN

Corrosive. Causes irreversible eye damage. Causes skin irritation. Harmful if inhaled. Harmful if swallowed. Harmful if absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic skin reactions in some individuals. Causes asthmatic signs and symptoms in hyper-reactive individuals.

Do not get in eyes, on skin, on clothing.  
Avoid breathing vapor. Do not swallow.  
Wear goggles, protective clothing, and butyl or nitrile gloves.  
Wash thoroughly with soap and water after handling.  
Remove contaminated clothing and wash before reuse.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

### STORAGE AND HANDLING

GLUTEX GS2 is incompatible with many commonly used materials of construction such as steel, galvanized iron, aluminum, tin, and zinc. GLUTEX GS2 can be stored and handled in baked phenolic-lined steel, polyethylene, stainless steel, or reinforced epoxy-plastic equipment. This product freezes at about 16°F (-9 °C). Therefore, unless the storage tank is inside or underground, heating and insulation may be required. If heating is needed, exposure to high temperatures should be avoided. For short storage times (up to about 1 month), temperatures of up to 37.8° C can be tolerated but the preferred maximum storage temperature is about (26.7° C).

A stainless steel centrifugal pump is suggested for transfer service. Spiral-wound stainless steel with TEFLON® Polymer is suitable for gaskets and packing.

Handle in a well-ventilated area. If vapors are irritating to the nose or eyes, special ventilation or respiratory protection (MSHA/NIOSH approved air purifying respirator equipped with an organic vapor cartridge) may be required.

### LIMITED WARRANTY AND DISCLAIMER


Seller warrants that the product conforms to its chemical description as contained on this label and is reasonably fit for the purposes stated on this label when used in accordance with directions under normal conditions of use. THE WARRANTIES MADE IN THIS PARAGRAPH ARE SELLER'S SOLE WARRANTIES WITH RESPECT TO THE PRODUCT AND ARE MADE EXPRESSLY IN LIEU OF AND EXCLUDE ANY IMPLIED WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE AND ALL OTHER EXPRESS OR IMPLIED REPRESENTATIONS AND WARRANTIES.

**BEFORE HANDLING OR USING THIS PRODUCT, SEE YOUR EMPLOYER AND READ CURRENT MATERIAL SAFETY DATA SHEET.**

## STORAGE AND DISPOSAL

**PESTICIDE DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or your Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL: Metal Containers or Plastic Containers:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or other procedures approved by state and local authorities. **Plastic Containers:** May be incinerated, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. **Metal Containers:** Must not be incinerated. Do not cut or weld on or near metal containers.



## NOTICE

Do Not Ship or Store with Food, Feeds, Drugs, or Clothing

85409-7/17/2006-965

# GLUTEX™ GS2

**AN INDUSTRIAL SANITIZER PRODUCT FOR USE IN CONTROLLING MICROBIAL CONTAMINATION IN ANIMAL PRODUCTION FACILITIES AND FARM EQUIPMENT SUCH AS: POULTRY & TURKEY HOUSES, SWINE HOUSING AND FARROWING AREAS, BARNs AND LARGE ANIMAL BUILDINGS, HATCHERS, SETTERS, AND CHICK PROCESSING FACILITIES, CAGES AND TRANSPORT VEHICLES, AND HATCHING EGGS AND FOR SANITIZING INDUSTRIAL OR NON-FOOD CONTACT SURFACES; SUCH AS EQUIPMENT FOR MIXING, PACKING, OR PACKAGING PRODUCTS, PRODUCT TRANSFER LINES, STORAGE TANKS, FLOORS AND WALLS.**

Active Ingredient: Glutaraldehyde 20%  
Inert Ingredients: 80%  
Total 100%

## KEEP OUT OF REACH OF CHILDREN DANGER

### FIRST AID

#### If swallowed:

- Call CHEMTREC or a doctor immediately for treatment advice.
- DO NOT INDUCE VOMITING.
- Do not give anything to drink.

#### If in eyes:

- Wash immediately and continuously with flowing water for at least 30 minutes.
- Remove contact lenses after the first 5 minutes and continue washing.
- Obtain prompt medical consultation, preferably from an ophthalmologist.

#### If on skin or clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call CHEMTREC or a doctor for treatment advice.

#### If inhaled:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, and then give artificial respiration, preferably mouth-to-mouth if possible.
- Call CHEMTREC or a doctor for further treatment advice.

**NOTE TO PHYSICIAN:** Aspiration may cause lung damage. Probable mucosal damage may contraindicate the use of gastric lavage.

Have the MSDS and, if available, the product container or label with you when calling a poison control center or a doctor, or going for treatment.

### SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS



### THE DOW CHEMICAL COMPANY

Midland, Michigan 48674 U.S.A.  
(989)636-4400

© TM Trademark of The Dow Chemical Company ("Dow")  
or an affiliated company of Dow

Made in U.S.A.

EPA Reg. No. 464-689  
EPA Est. No. 10352-WV-2 (A); 464-WV-1 (B)  
EPA Est. No. 10352-WV-1 (C)  
(A) 5 gallon pails in pallets; all drums; bulk  
(B) Single 5 gallon pails only.  
(C) Bulk Shipments

Net Contents: 5 or 55 gallons

CONTAINER SIZE 5 GALLONS	CONTAINER SIZE 55 GALLONS
43 POUNDS NET WEIGHT	470 POUNDS NET WEIGHT

L55911 (107/04)  
Made In USA  
PE0506

**IN CASE OF AN EMERGENCY** endangering life or property involving this product, call collect (989)-636-4400.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

### SANITIZING NON-FOOD CONTACT SURFACES, FARM, ANIMAL, AND POULTRY HOUSING FACILITIES AND EQUIPMENT

#### DO NOT USE THIS PRODUCT ON FOOD OR FEED

Product dilution:

To **sanitize vegetative pathogens**, *Clostridium perfringens*, *Enterobacter aerogenes*, *Escherichia coli*, *Haemophilus parasuis*, *Klebsiella pneumoniae*, *Mycoplasma gallisepticum*, *Mycoplasma synoviae*, *Pasteurella multocida*, *Salmonella enteritidis*, *Salmonella pullorum*, *Salmonella typhi*, *Staphylococcus aureus*, and *Streptococcus suis*, **and viruses:** *Avian infectious bronchitis virus*, *Avian influenza*, *Avian laryngotracheitis*, *Avian reovirus*, *Avian rotavirus*, *Canine parvovirus*, *Infectious bursal disease*, *Marek's disease virus*, *Pseudorabies virus*, *Transmissible gastroenteritis virus*, *Hog Cholera virus*, *Porcine reproductive respiratory syndrome* and *Human Coronavirus*.

1. Thoroughly saturate all surfaces using a mop or by spraying with a solution prepared by mixing 0.61 to 1.5 fluid ounces of GLUTEX GS2 per gallon of water to produce a solution containing 0.1 to 0.25% active ingredient.
2. Allow to stand for at least five minutes, or until thoroughly dried.
3. Ventilate thoroughly before re-use.
4. Fresh solution should be prepared daily or more often if the solution becomes diluted or soiled.

#### Farm Equipment and Animal Housing Buildings

(poultry & turkey grow-out houses, laying houses, swine production and housing, barns and large animal buildings)

Before applying GLUTEX GS2:

1. Remove all poultry and other animals and feed from the premises. Remove all portable equipment and empty all feeders and water troughs. If not removed, all treated feeders and waterers must be thoroughly rinsed with potable water prior to re-use.
2. Remove all litter and manure from floors and other surfaces.
3. Using suitable commercially available equipment, saturate all surfaces with diluted GLUTEX GS2 (see Product Dilution, above).
4. Allow to stand for at least five minutes or until dried.
5. Ventilate enclosed spaces.
6. Do not repopulate with poultry or other animals, or use equipment until treatment has been absorbed or dried.

#### Hatchers, Setters, and Chick Processing Facilities

1. General sanitizing of environmental surfaces prior to introduction of eggs: Remove all animals from the area. Clean out feathers, fluff, or other debris. Thoroughly saturate all surfaces with a solution prepared by mixing the appropriate amount (see Product Dilution, above) of GLUTEX GS2 with water. Allow to stand for at least five minutes or until completely dried. Ventilate thoroughly before reuse.
2. Treatment of air and surfaces after introduction of eggs in high humidity hatchers and setters with water spray cooling: Prepare a solution by mixing the appropriate amount of GLUTEX GS2 with water (see Product Dilution, above). Fog or atomize this solution into setters or hatchers, using appropriate application equipment.
3. Treatment of air and surfaces after introduction of eggs in low humidity hatchers with chilled coil cooling: Fog or atomize this product using appropriate application equipment.

#### Trays, Racks, Carts, Chick Boxes, Cages, and other Hard Surfaces:

1. Remove all filth and heavy debris from surfaces by scraping or washing.
2. Prepare a solution by mixing the appropriate amount (see Product Dilution, above) of GLUTEX GS2 with water.
3. Apply by spraying or immersion, and allow to drain and air-dry.

#### Trucks and other Vehicles

1. Remove all filth and heavy debris from surfaces by scraping or washing.
2. Prepare a solution by mixing the appropriate amount (see Product Dilution, above) of GLUTEX GS2 with water.
3. Apply by spraying or immersion, and allow to air-dry.

#### Egg Sanitization

(for treating hatching eggs only)

Prepare a solution by mixing 0.64 ounces of GLUTEX GS2 with one gallon of water. Place in a bath at 105° to 115°F (40° to 45°C). Place freshly laid eggs in plastic trays and dip briefly (10 to 15 seconds), then allow to drain and air dry.

### SANITIZING NON-FOOD CONTACT SURFACES, INDUSTRIAL EQUIPMENT AND BUILDINGS

Before applying GLUTEX GS2, remove heavy filth and soil deposits from surfaces using a suitable cleaner. Dilute to 0.10% to 0.25% active ingredient by mixing 0.61 to 1.5 fluid ounces of GLUTEX GS2 per gallon of water. Using suitable commercially available equipment, saturate all surfaces with diluted GLUTEX GS2. Allow to set for at least five minutes or until dried. Ventilate enclosed spaces.