

INTERVET/SCHERING-PLOUGH ANIMAL HEALTH (formerly Intervet Inc.)

29160 INTERVET LANE, P.O. BOX 318, MILLSBORO, DE, 19966-0318

Toll-Free: 800-992-8051
 Customer Service: 800-441-8272
 Website: www.intervetusa.com
 Email: Information.USA@intervet.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the US product label or package insert.

NEWHATCH-C2-M™

Intervet

Newcastle-Bronchitis Vaccine

B1 Type, C2 Strain, Mass. Type

Live Virus

Coarse Spray

READ IN FULL

DESCRIPTION

NEWHATCH-C2-M, a live virus vaccine, is prepared from a B₁ Type, C2 Strain of Newcastle disease virus and a mild Massachusetts strain of Infectious Bronchitis virus. The viruses have been propagated using SPF substrates.

INDICATIONS FOR USE

The vaccine is recommended for vaccination of healthy chickens, one day of age or older for protection against Newcastle disease and Infectious Bronchitis disease (Massachusetts strain) by coarse spray administration.

VACCINATION PROGRAM

Many factors must be considered in determining a sound vaccination program for a particular farm or poultry complex. To be fully effective, the vaccine must be administered properly to healthy, receptive chickens maintained in a proper environment under good management. In addition, the response may be influenced by the age of the chickens and their immune status. Seldom does one live virus vaccination under field conditions produce lifetime protection for all individuals in a given flock. The level of immunity required will vary with operational practices and the degree of exposure. Therefore, a program of periodic revaccinations may be necessary.

PREPARATION OF VACCINE

For Coarse Spray

Product Code NB-208-10

Do not open and mix the vaccine until ready to begin vaccination. Use vaccine immediately after mixing.

1. Remove the tear-off seal and stopper from the vial containing the dried vaccine.
2. Carefully pour clean, cool, deionized water into the vaccine vial until the vial is approximately two-thirds full.
3. Insert the rubber stopper and shake vigorously until all material is dissolved.
4. The vaccine is now ready for coarse spray use in accordance with the following directions. For best results, be sure to follow directions carefully!

COARSE SPRAY ADMINISTRATION - For Chickens One Day of Age

1. Use rehydrated vaccine as indicated for specific coarse spray vaccination machine. For example, a machine which dispenses 20 ml to a box of 100 chickens - total volume for 10,000 doses is 2,000 ml of deionized water. Mix thoroughly.
2. Add the prepared vaccine solution to reservoir on the machine.
3. Prime and adjust machine as instructed in manual accompanying the specific machine.
4. Place boxes holding 100 chickens each on the conveyor belt or in machine. Activate spray head.

CAUTION

1. VACCINATE ONLY HEALTHY CHICKENS. Although disease may not be evident, Coccidiosis, Mycoplasma infection, Infectious Bursal disease, Chicken Infectious Anemia, Reovirus infection, Marek's disease and other disease conditions may cause complications or reduce immunity.
2. All susceptible chickens on the same premises should be vaccinated at the same time.
3. The revaccination of laying hens with live Newcastle/Bronchitis vaccine may be detrimental to the flock and

cannot be generally recommended. This caution applies to all live Newcastle/Bronchitis vaccines currently available. Consult your Intervet representative for more information.

4. Efforts should be taken to reduce stress conditions at the time of vaccination and during the reaction period.
5. Do not spill or splash the vaccine.
6. Do not dilute the vaccine or otherwise stretch the dosage.
7. Use entire contents when first opened.
8. Do not vaccinate within 21 days before slaughter.
9. Burn containers and all unused contents.
10. Avoid exposure of this vaccine to sunlight.
11. Do not use less than 1 dose per bird.
12. This vaccine contains gentamicin as a preservative.
13. **WARNING:** Newcastle virus occasionally causes conjunctivitis in humans. Avoid any contact of vaccine with eyes.
14. For veterinary use only.

NOTICE

This vaccine has undergone rigid potency, safety and purity tests and meets Intervet Inc. and USDA requirements. It is designed to stimulate effective immunity when used as directed, but the user must be advised that the response to the product depends upon many factors, including, but not limited to, conditions of storage and handling by the user, administration of the vaccine, health and responsiveness of individual chickens and the degree of field exposure. Therefore, directions should be followed carefully.

This product is not hazardous when used according to directions supplied. A material safety data sheet (MSDS) is available upon request. This and any other consumer information can be obtained by calling Intervet Customer Service at 1-800-441-8272 or 1-302-934-8051.

RECORDS

Keep a record of vaccine, quantity, serial number, expiration date and place of purchase; the date and time of vaccination; the number, age, breed and locations of chickens; names of operators performing the vaccination and any observed reactions.

HOW SUPPLIED

		Ordering Number
10 X 10,000 Doses	Coarse Spray Use	NB-208-10

The use of this vaccine is subject to applicable federal and local laws and regulations.

Use only as directed

Store vaccine between 2 and 7°C (35 and 45°F)

This product is nonreturnable

INTERVET INC.

Millsboro, Delaware U.S.A.

U.S. Veterinary License No. 286

Made in U.S.A.

U.S. Patent No. 5,750,111

20803

AL176

NAC No.: 11062830