

Oxytet™

Soluble

Oxytetracycline hydrochloride

Soluble powder for use in the drinking water

- Broad-spectrum antibiotic
- Citric acid provides maximum absorption
- Excellent solubility
- Superior palatability
- 3 packet sizes to meet wide variety of flock or herd sizes

Composition:

- Oxytetracycline hydrochloride antibiotic powder
- 70 gm packet (25.6 active grams)
 - 280 gm packet (102.4 active grams)
 - 1400 gm packet (512.0 active grams)

Indications:

- Chickens - for the control of infectious synovitis (*Mycoplasma synoviae*/MS) and CRD (chronic respiratory disease) and air sac infection (*M. gallisepticum*/MG); also for the control of colibacillosis (*E. coli*) and Fowl cholera (*Pasteurella multocida*)
- Growing turkeys - for the control of complicating bacterial organisms associated with bluecomb (transmissible enteritis/coronaviral enteritis)
- Turkeys – for the control of hexamitiasis (*Hexamita meleagridis*) and infectious synovitis (*M. synoviae*)
- Swine – to control and treat bacterial enteritis (*E. coli* and *Salmonella choleraesuis*) and bacterial pneumonia (*P. multocida*)
- Breeding swine – to control and treat leptospirosis (*Leptospira pomona*) and thereby reduce the incidence of abortions and shedding of spirochetes

**All of the above organisms must be susceptible to tetracycline for effective treatment.*

For more information on putting Oxytet Soluble to work for you, contact your nearest Alpha representative.

Alpha – The performance enhancement people

Oxytet®

Soluble



AlphaPharma Inc.
One Executive Drive
Fort Lee, NJ 07024, USA

Tel (201) 947-7774
Fax (201) 585-0429
<http://www.alphaPharma.com>

Oxytet is a trademark of AlphaPharma Inc.

Copyright ©2001 by AlphaPharma Inc.

Proportioner Mixing Instructions:

Add the following amount of Oxytet Soluble to one gallon of water to make a stock solution for proportioner set to meter at the rate of one ounce per gallon.

Species	Disease	Treatment Level	Grams of Product / Gallon Stock Solution	Withdrawal	Treatment
Chicken ¹	Infection synovitis (MS)	200 - 400 mg/gal.	70 - 140 grams	0 days	7 - 14 days
	Chronic Respiratory Disease (MG) or <i>E. coli</i>	400 - 800 mg/gal.	140 - 280 grams	0 days	7 - 14 days
	Fowl Cholera (<i>P. multocida</i>)	400 - 800 mg/gal.	140 - 280 grams	0 days	7 - 14 days
	Hexamitiasis	200 - 400 mg/gal.	70 - 140 grams	0 days	7 - 14 days
Turkey ¹	Infectious synovitis (MS)	400 mg/gal.	140 grams	0 days	7 - 14 days
	Coronaviral enteritis Secondary bacterial infection	25 mg/lb of body weight	----	0 days	7 - 14 days
Swine	Bacterial enteritis (<i>E. Coli</i> , <i>Salmonella choleraesuis</i>)	10 mg/lb of body weight	----	0 days	1 - 5 days
	Bacterial pneumonia (<i>P. multocida</i>)	10 mg/lb of body weight	----	0 days	1 - 5 days
Breeding Swine	Leptospirosis (<i>L. pomona</i>)	10 mg/lb of body weight	----	0 days	1 - 5 days

¹Do not use in birds producing eggs for human consumption.

Tank Mixing Instructions:

Each package size when mixed with the volumes of water listed below makes treatment solutions with the corresponding oxytetracycline HCl concentrations. See above for indications and use levels.

Solution Levels	Amount of Product		
	70 grams	280 grams	1400 grams
200 mg solution	128 gal. water	128 gal. water	2560 gal. water
400 mg solution	64 gal. water	64 gal. water	1280 gal. water
800 mg solution	32 gal. water	32 gal. water	640 gal. water

See package label for complete use directions and warnings.

Package sizes:

- 100 x 70 gram
- 50 x 280 gram
- 10 x 1400 gram