

# SpecLinx-50<sup>®</sup>

(Brand of Lincomycin-Spectinomycin soluble powder)  
Water Soluble Powder

ANADA 200-380, Approved by FDA



## INDICATIONS

For use in chickens up to 7 days of age as an aid in the control of: Airsacculitis caused by either *Mycoplasma synoviae* or *Mycoplasma gallisepticum* susceptible to lincomycin-spectinomycin.

Complicated Chronic Respiratory Disease (Air Sac Infection) caused by *Escherichia coli* and *M. gallisepticum* susceptible to lincomycin-spectinomycin.

See reverse side for Administration and Dosage.

## BENEFITS

- **Highly soluble:** Mixes easily and completely in water
- **Economical:** A single 375 g packet treats 125 gallons of water
- **Convenient:** Pre-measured packaging provides for handy on-site mixing
- **Stable:** Stores well at room temperature
- **FDA approved**

## PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1LIN009	75 g pouch	12
1LIN010	375 g pouch	6

# SpecLinx-50<sup>®</sup>

(Brand of Lincomycin-Spectinomycin soluble powder)  
Water Soluble Powder

ANADA 200-380, Approved by FDA

### INGREDIENTS

Each 75 g of SpecLinx-50 contains:  
 Lincomycin hydrochloride  
 Equivalent to lincomycin ..... 16.7 grams  
 Spectinomycin dihydrochloride pentahydrate  
 Equivalent to spectinomycin ..... 33.3 grams  
 Total Antibiotic Activity ..... 50.0 grams

### INDICATIONS

For use in chickens up to 7 days of age as an aid in the control of: *Airsacculitis* caused by either *Mycoplasma synoviae* or *Mycoplasma gallisepticum* susceptible to lincomycin-spectinomycin.  
 Complicated Chronic Respiratory Disease (Air Sac Infection) caused by *Escherichia coli* and *M. gallisepticum* susceptible to lincomycin-spectinomycin.

### DOSAGE

Provide 2 grams (g) antibiotic activity per gallon of drinking water. Administer as the sole source of water for the first 5 to 7 days of life.

Administration	
Amount of Spec-Linx 50	Amount of Drinking Water
75 g	25 Gallons
375 g	125 Gallons

For proportioners delivering 1 oz/gallon of drinking water, dissolve the contents of 5 packets of the 75 g pouch or 1 packet of the 375 g pouch size in each gallon of proportioner solution.

### IMPORTANT

Chickens should consume water at the following approximate rate to insure intake of the required dose of lincomycin-spectinomycin indicated:

Broilers and Layer Replacements (Light and Heavy)		
Age (Weeks)	Daily Water intake Gallons/1000	Dosage mg Antibiotic/lb
1	5	50-65

### CAUTION

Discard medicated drinking water daily and replace with fresh medicated drinking water. Store at controlled room temperature 20° to 25°C (68° to 77°F).

### WARNING

**NOT FOR HUMAN USE**  
**KEEP OUT OF REACH OF CHILDREN**

### HOW SUPPLIED

SpecLinx-50 is supplied in 75 g (2.65 oz) and 375 g (13.2 oz) foil pouches; and 75 g (2.65 oz) plastic jars.



SpecLinx is a Registered Trademark of Bimeda, Inc.