

Valbazen®
(albendazole)

Broad-Spectrum Dewormer

Oral Suspension for Use in Cattle and Sheep

Controls:

Stomach Worms

including 4th stage inhibited larvae of *Ostertagia ostertagi*

Intestinal Worms

Lungworms

Tapeworms

Mature Liver Fluke

For removal and control of liver flukes, tapeworms, stomach worms, intestinal worms, lungworms

Active Ingredient:

Albendazole 11.36%
(Equivalent to 113.6 mg/mL)

Indications: Valbazen is a broad-spectrum anthelmintic effective in the removal and control of the following internal parasites in cattle and sheep:

Parasite	Cattle	Sheep
Adult Liver Flukes	<i>Fasciola hepatica</i>	<i>Fasciola hepatica, Fascioloides magna</i>
Heads and Segments of Tapeworms	<i>Moniezia benedeni, M. expansa</i>	Common Tapeworms (<i>Moniezia expansa</i>) Fringed Tapeworms (<i>Thysanosoma actinioides</i>)
Adult and 4th Stage Larvae of Stomach Worms	Brown Stomach Worms, including 4th stage inhibited larvae (<i>Ostertagia ostertagi</i>), Barberpole Worm (<i>Haemonchus contortus, H. placei</i>), Small Stomach Worm (<i>Trichostrongylus axei</i>)	Brown Stomach Worms (<i>Ostertagia circumcincta, Marshallagia marshalli</i>), Barberpole Worm (<i>Haemonchus contortus</i>), Small Stomach Worm (<i>Trichostrongylus axei</i>)
Adult and 4th Stage Larvae of Intestinal Worms	Thread-necked Intestinal Worm (<i>Nematodirus spathiger, N. helvetianus</i>), Small Intestinal Worm (<i>Cooperia punctata, C. oncophora</i>)	Thread-necked Intestinal Worm (<i>Nematodirus spathiger, N. filicollis</i>), Cooper's Worms (<i>Cooperia oncophora</i>), Bankrupt Worm (<i>Trichostrongylus colubriformis</i>), Nodular Worm (<i>Oesophagostomum columbianum</i>), Large-mouth Bowel Worm (<i>Chabertia ovina</i>)
Adult Stages of Intestinal Worms	Hookworm (<i>Bunostomum phlebotomum</i>), Bankrupt Worm (<i>Trichostrongylus colubriformis</i>), Nodular Worm (<i>Oesophagostomum radiatum</i>)	
Adult and 4th Stage Larvae of Lungworms		<i>Dictyocaulus viviparus</i>
Adult and Larval Stages of Lungworms		<i>Dictyocaulus filaria</i>

Dosage and Administration: Valbazen Suspension should be administered to cattle at the recommended rate of 4 mL/100 lb of body weight (equivalent to 4.54 mg of albendazole/lb, 10 mg/kg) and to sheep at the recommended rate of 0.75 mL/25 lb of body weight (equivalent to 3.4 mg of albendazole/lb, 7.5 mg/kg). The following table indicates recommended dosing schedules.

Cattle		Sheep	
Body Weight	Dosage	Body Weight	Dosage
250 lb	10 mL	25 lb	0.75 mL
500 lb	20 mL	50 lb	1.5 mL
750 lb	30 mL	75 lb	2.25 mL
1000 lb	40 mL	100 lb	3.0 mL
1250 lb	50 mL	200 lb	6.0 mL
1500 lb	60 mL	300 lb	9.0 mL

Cattle: 5 liters of Valbazen 11.36% Suspension will treat 250 animals weighing 500 lb.

Sheep: 5 liters of Valbazen 11.36% Suspension will treat 3320 animals weighing 50 lb.

Valbazen 11.36% Suspension should be given orally using any type of standard dosing gun or dose syringe.

Important: Accurate estimates of the weight of the cattle and sheep to be treated are essential for most effective results with this product. Animals constantly exposed to internal parasites should be retreated as necessary.

Warnings: Cattle must not be slaughtered within 27 days following last treatment. Sheep must not be slaughtered within 7 days following last treatment. Because a withdrawal time in milk has not been established, do not use in female dairy cattle of breeding age.

Caution: Do not administer to female cattle during first 45 days of pregnancy or for 45 days after removal of bulls. Do not administer to ewes during the first 30 days of pregnancy or for 30 days after removal of rams. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

Keep This and All Medication Out of Reach of Children

Shake Well Before Using

For Use in Animals Only

Store at Controlled Room Temperature 15°–30°C (59°–86°F)

Protect From Freezing

U.S. Patent Nos. 3,915,986 and 3,956,499

NADA #110-048, Approved by FDA

Distributed by:

Pfizer Animal Health

Exton, PA 19341, USA

Div. of Pfizer Inc

NY, NY 10017

85-8785-05