Lincomycin

Soluble

Lincomycin hydrochloride soluble powder

Drinking water administration

- Water-soluble antibiotic with gram-positive spectrum of activity
- Two convenient package sizes for easy use and accurate administration to broiler chickens or swine
- Immediate treatment for important enteric disease of chickens and swine

Composition:

Lincomycin Soluble is a water-soluble form of the antibiotic lincomycin.

Indications/Dose:

- Lincomycin Soluble is for use in swine for the treatment of swine dysentery (bloody scours).
- In swine, **Lincomycin Soluble** should be administered for a minimum of 5 consecutive days beyond the disappearance of symptoms (bloody feces) up to a maximum of 10 consecutive days.
- In chickens, **Lincomycin Soluble** is for the control of necrotic enteritis caused by *Clostridium perfringens* susceptible to lincomycin.
- In chickens, **Lincomycin Soluble** should be administered for 7 consecutive days.

For more information on putting Lincomycin Soluble to work for you, contact your nearest Alpharma representative.

Alpharma – The performance enhancement people

Lincomycin Soluble



Alpharma Inc. One Executive Drive Fort Lee, NJ 07024, USA Tel (201) 947-7774 Fax (201) 585-0429 http://www.alpharma.com



Proportioner Mixing Instructions:

Add the appropriate package (40 gram or 80 gram) of Lincomycin Soluble to water to make a stock solution for proportioner set to meter at a rate of one ounce per gallon (1:128).

Species	Disease	Stock Solution from 40 gm packet	Stock Solution from 80 gm packet	Withdrawal
Chicken ¹	Necrotic enteritis	1.95 gallons	3.9 gallons	0 Days
Swine ²	Swine dysentery Dose: 3.8mg/pound/day	0.5 gallons	1.0 gallons	6 Days

Tank Mixing Instructions:

Add the appropriate package (40 gram or 80 gram) of Lincomycin Soluble to a tank of adequate volume and then add the appropriate amount of water to make the treatment solutions listed below.

Species	Disease	Treatment Solution from 40 gm packet	Treatment Solution from 80 gm packet	Withdrawal
Chicken ¹	Necrotic enteritis	250 gallons	500 gallons	0 Days
Swine ²	Swine dysentery Dose: 3.8mg/pound/day	64 gallons	128 gallons	6 Days

¹ Not for use in layers and breeder chickens.

² Do not use for more than 10 days. Not for use in swine weighing more than 250 pounds.

See package label for complete use directions and warnings.

Package size:

25 x 40 grams 25 x 80 grams