

NeoMed[®] 325

(neomycin sulfate) Soluble Powder

ANADA 200-050, Approved by FDA



INDICATIONS

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in cattle (excluding veal calves), swine, sheep and goats. For the control of mortality associated with *Escherichia coli* organisms susceptible to neomycin sulfate in growing turkeys.

PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1NE0013	100 gm	50
1NE0010	200 gm	25
1NE0007	50 lb.	1

BENEFITS

- **Safe:** Approved for treatment of growing turkeys with no withdrawal period
- **Versatile:** Provides treatment for a wide variety of species
- **Soluble:** Dissolves readily in drinking water
- **Convenient:** Available in 100 g and 200 g packets plus space saving resealable 50 lb. bulk pails
- **FDA approved**

See reverse side for Administration and Dosage.

NeoMed[®]-325

(neomycin sulfate) Soluble Powder

ANADA 200-050, Approved by FDA

DESCRIPTION

Add to drinking water - Not for use in liquid supplements.

INDICATIONS

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in cattle (excluding veal calves), swine, sheep and goats. For the control of mortality associated with *Escherichia coli* organisms susceptible to neomycin sulfate in growing turkeys.

DOSAGE & ADMINISTRATION

Administer to cattle (excluding veal calves), swine, sheep and goats at a dose of 10 mg neomycin sulfate per pound of body weight per day in divided doses for a maximum of 14 days. Administer to turkeys at a dose of 10 mg neomycin sulfate per pound of body weight per day for 5 days.

HERD/FLOCK TREATMENT

100 g (3.5 oz) NeoMed 325 will treat 7150 pounds of body weight. Estimate the total number of pounds of body weight of the animals to be treated and administer 3.5 oz for each 7150 pounds. The product should be added to the amount of drinking water estimated to be consumed in 12-24 hours. Provide medicated water as the sole source of water each day until consumed, followed by non-medicated water as required. Fresh medicated water should be prepared each day.

INDIVIDUAL ANIMAL TREATMENT

To provide 10 mg neomycin sulfate per pound of body weight, mix one (1) level teaspoon in water or milk for each 160 pounds body weight. Administer daily either as a drench in divided doses or in the drinking water to be consumed in 12-24 hours.

DRINKING WATER

Swine - Use the amount of NeoMed 325 indicated below in 256 gallons of water, or in two gallons of stock solution used in proportioner set to meter one ounce per gallon.

Pigs weighing 25 to 50 Pounds.....	7 oz
Pigs weighing 50 to 100 Pounds.....	10.5 oz
Pigs weighing over 100 Pounds.....	14 oz

CAUTION

To administer the stated dosage, the concentration of neomycin required in medicated water must be adjusted to compensate for variation in age and weight of animal, the nature and severity of disease signs, and environmental temperature and humidity, each of which affects water consumption.

If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. If symptoms such as fever, depression, or going off feed develop, oral neomycin is not indicated as the sole treatment since systemic levels of neomycin are not obtained due to low absorption from the gastrointestinal tract.

IMPORTANT

Treatment should continue 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 14 consecutive days for cattle, swine, sheep and goats and 5 days for turkeys. Animals not drinking or eating should be treated individually by drench.

WARNING

DISCONTINUE TREATMENT PRIOR TO SLAUGHTER BY AT LEAST THE NUMBER OF DAYS LISTED BELOW FOR APPROPRIATE SPECIES:

Cattle.....	1 day
Swine and goats.....	3 days
Sheep.....	2 days
Turkeys.....	0 days

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older.

Restricted Drug (California), use only as directed.

Securely reseal bulk pails and tightly fold packets to keep out moisture.

Store at controlled room temperature.

HOW SUPPLIED

NeoMed 325 is supplied in 100 g (3.5 oz) pouch, packed 50 per pail; 200 g (7.0 oz) pouch packed 25 per pail; and 50 lb. resealable bulk pail.

TAKE TIME



OBSERVE LABEL
DIRECTIONS