

Penicillin G Potassium USP

For Animal Use Only

Not for Human Use

ANTIBIOTIC FOR DRINKING WATER

FOR ORAL USE IN TURKEYS ONLY

NDC 59320-001-02

ACTIVE INGREDIENT:

0.500 billion units Penicillin G Potassium

Nonsterile

KEEP OUT OF REACH OF CHILDREN

Restricted Drug (California) — Use Only as Directed

INDICATIONS: For treatment of erysipelas in turkeys caused by *Erysipelothrix rhusiopathiae*.

RESIDUE WARNINGS: Treated turkeys must not be slaughtered for food during treatment and for one day after last treatment.

Do not use in turkeys producing eggs for human consumption.

PRECAUTIONS: For best results, the treatment should be started at the first sign of infection. If improvement is not noted after 3 to 4 days of treatment, consult a poultry pathologist or veterinarian. Keep this and all medication out of reach of children.

STORAGE AND HANDLING: Store at or below 25°C (77°F). Protect from excessive heat, 40°C (104°F), and moisture. Label recommendations for storage and replacement of stock and medicated water solutions must be followed to assure the performance of this drug product.

DOSAGE AND ADMINISTRATION:

Administer orally at a dosage of 1,500,000 units of penicillin per gallon (3.8 liters) of drinking water for 5 consecutive days.

DIRECTIONS: Combine contents and approximately 2 pints (946 mL) of water in glass or plastic container. Stir to dissolve. Allow the concentrated solution to stand until the foam disappears. The concentrated solution should be used up or discarded within one hour of preparation.

Automatic Watering Systems: Pour the concentrated solution into a glass or plastic container then add enough water to make 2.6 gallons (9.9 liters) of stock solution. {This amount of solution will medicate 333 gallons (1260 liters) of

drinking water}. The automatic waterer should be adjusted to deliver 1 oz (30 mL) of stock solution per gallon (3.8 liters) of drinking water. Prepare fresh stock solutions and medicated drinking water solutions every twelve (12) hours. All solutions in contact with galvanized metal should be changed every three (3) hours.

Gravity Flow Watering Systems: Pour the concentrated solution into enough water to make 333 gallons (1260 liters) of drinking water. In gravity flow watering systems, prepare fresh solution every 12 hours. All solutions in contact with galvanized metal should be changed every three (3) hours.

Drinking water prepared as directed above will contain 1,500,000 units of Penicillin G Potassium per gallon (3.8 liters).

ANADA 200-307, Approved by FDA

25148A

Made in Canada

RevDate: February 2010

Manufactured for:

Vetoquinol N.-A. Inc.

2000, ch. Georges,

Lavaltrie, QC, Canada J5T 3S5

Penicillin G Potassium USP

For Animal Use Only

Not for Human Use

ANTIBIOTIC FOR DRINKING WATER

FOR ORAL USE IN TURKEYS ONLY

NDC 59320-001-02

ACTIVE INGREDIENT:

0.500 billion units Penicillin G Potassium

Nonsterile

KEEP OUT OF REACH OF CHILDREN

Restricted Drug (California) — Use Only as Directed