

# Di-Methox<sup>®</sup>

## Sulfadimethoxine 12.5% Oral Solution

AgriLabs offers a wide variety of effective pharmaceutical products, including **Di-Methox** Oral Solution which is the product of choice because of its broad coverage of a variety of diseases. **Di-Methox** Oral Solution has increased in popularity because it is considered highly effective in the treatment of pneumonia, shipping fever, calf diphtheria, and foot rot in cattle. It also treats outbreaks of coccidiosis and fowl cholera. **Di-Methox** Oral Solution is cost-effective, safe and the product of choice for use in your operation.

- ✓ Broad coverage for cattle
  - Shipping Fever
  - Pneumonia
  - Calf Diphtheria
  - Foot Rot
- ✓ Coccidiosis and fowl cholera
- ✓ Low stress delivery system
- ✓ Oral dosing
- ✓ Cost Effective and Safe
- ✓ High Quality and Proven Product



Technical Support is only a phone call away.  
800-542-8916

**Di-Methox<sup>®</sup>**  
**12.5% Oral Solution**

**INDICATIONS:**  
**CHICKENS AND TURKEYS:** Use for the treatment of disease outbreaks of coccidiosis, foot cholera, and salmonellosis.

**MEAT PRODUCTION THERAPY:** Use for the treatment of disease outbreaks of coccidiosis and foot cholera in meat chickens, turkey broilers and turkey culls. Use in the treatment of shipping fever complex, bacterial pneumonia, and other enteric and head end.

**CAUTION:** Store at room temperature. If freezing occurs, freeze before using. Protect from light. Direct sunlight may cause discoloration. Protecting an administration does not affect potency. Prepare a fresh stock solution daily.

**CHICKENS AND TURKEYS:** If animals show the signs of disease within 5 days, coccidiosis or bacterial enteric or respiratory diagnosis. Follow the recommended dosage (chickens water and turkey 1/2 oz) or a cup per 100 lbs of water. Administer in one dose source of drinking water and sulfonamide medication, 12 hours, and the days that birds consumed feed. Administer a solution that should not be used for other purposes or feeders.

**CATTLE:** During treatment period, make certain that animals receive adequate water. Administer 1/2 oz per 100 lbs of water movement within 2 or 3 days. Re-evaluate diagnosis. Treat until animals are free of clinical signs for 3 days.

**WARNING:**  
**CHICKENS AND TURKEYS:** Administer 5 days before slaughter. Do not administer to animals less than 10 weeks (110 days) of age or to turkeys under 16 weeks of age.  
**CATTLE:** Withhold 7 days before slaughter. For dairy calves, dairy heifers, and adult cattle only. A withdrawal period for milk residues is indicated for use, pending a pending withdrawal. DO NOT USE IN OTHER THAN INDICATED PURPOSES.


**FOR USE IN DRINKING WATER ONLY**  
 Do not use in feed or water (not in drinking water only as directed)  
 NOT FOR HUMAN USE  
 MULTIPLE USE IN CHICKENS AND TURKEYS

Manufactured for:  
 Agri Laboratories, Ltd.  
 St. Joseph, MO 64503

LOT NO.                      EXP. DATE

500014

# DI-METHOX<sup>®</sup>



## SULFADIMETHOXINE 12.5% ORAL SOLUTION ANTIBACTERIAL

For Oral Use in Chickens,  
 Turkeys and Cattle

12.5% Concentrated Solution  
 for Use in Drinking Water

ANADA 200-192, Approved by FDA

Net Contents: 3.785 L (1 GALLON) (128 fl oz)



**EACH FLUID OUNCE CONTAINS 375 mg SULFADIMETHOXINE SOLUBILIZED WITH ACACIA HYDRICOLA.**

Species	Concentration	Use Directions
CHICKENS	12.5%	1/2 oz per 100 lbs of body weight (1/2 cup per 20 lbs) in 20 gallons of drinking water
TURKEYS	12.5%	1/2 oz per 100 lbs of body weight (1/2 cup per 20 lbs) in 20 gallons of drinking water

**ADMINISTRATIVE:** Stock solution = 1/2 ounce per gallon of water.

Species	Concentration	Use Directions
CATTLE	12.5%	1/2 oz per 100 lbs of body weight (1/2 cup per 20 lbs) in 20 gallons of drinking water
TURKEYS	12.5%	1/2 oz per 100 lbs of body weight (1/2 cup per 20 lbs) in 20 gallons of drinking water

**TREATMENT PERIOD:** 5 consecutive days.

Species	Concentration	Use Directions
CHICKENS	12.5%	1/2 oz per 100 lbs of body weight (1/2 cup per 20 lbs) in 20 gallons of drinking water
TURKEYS	12.5%	1/2 oz per 100 lbs of body weight (1/2 cup per 20 lbs) in 20 gallons of drinking water
CATTLE	12.5%	1/2 oz per 100 lbs of body weight (1/2 cup per 20 lbs) in 20 gallons of drinking water
TURKEYS	12.5%	1/2 oz per 100 lbs of body weight (1/2 cup per 20 lbs) in 20 gallons of drinking water

**NOTE:** Withdraw 7 days before slaughter. For dairy calves, dairy heifers, and adult cattle only. A withdrawal period for milk residues is indicated for use, pending a pending withdrawal. DO NOT USE IN OTHER THAN INDICATED PURPOSES.

**WARNING:** CHICKENS AND TURKEYS: Administer 5 days before slaughter. Do not administer to animals less than 10 weeks (110 days) of age or to turkeys under 16 weeks of age. CATTLE: Withhold 7 days before slaughter. For dairy calves, dairy heifers, and adult cattle only. A withdrawal period for milk residues is indicated for use, pending a pending withdrawal. DO NOT USE IN OTHER THAN INDICATED PURPOSES.

**FOR USE IN DRINKING WATER ONLY**  
 Do not use in feed or water (not in drinking water only as directed)  
 NOT FOR HUMAN USE  
 MULTIPLE USE IN CHICKENS AND TURKEYS

Manufactured for:  
 Agri Laboratories, Ltd.  
 St. Joseph, MO 64503

LOT NO.                      EXP. DATE

500014



**Technical Support is only a phone call away.**  
**800-542-8916**

When building the technical service department, AgriLabs realized the need to hire veterinarians with practical field experience who also had industry expertise in cow/calf, dairy, and feedlot production medicine. AgriLabs veterinarians have hands-on practical knowledge, as well as many years of experience with product development, support trials, field investigations, and technical research projects. Our technical support staff provides producers with a knowledgeable friend in the business.

